



Informed Care Strategies

[LOG OUT](#) | [SEARCH](#) | [MY PRODUCTS](#) | [CONTACT US](#) | [USER GUIDE](#)

 27th Edition [AC](#) [ISC](#) [GRG](#) [MCM](#) [RFC](#) [HC](#) [CCG](#) [TC](#) [BHG](#) PIP [MCR](#) **Date of Service**

 Quick Search
[Recovery Facility Care](#) > [Optimal Recovery Guidelines](#) > [Subacute/Skilled Nursing Facility \(SNF\)](#) > [Cardiovascular Surgery](#)
 >[Aortofemoral Bypass \(S-5160\)](#)

Aortofemoral Bypass

ORG: S-5160 (RFC)

[Link to Codes](#)
MCG Health
 Recovery Facility
 Care
 27th Edition

- [Clinical Indications for Admission to Recovery Facility](#)
- [Alternative Levels of Care](#)
- [Length of Stay](#)
- [Evaluation and Treatment](#)
 - [General Treatment Course](#)
 - [Extended Stay](#)
 - [Patient Education](#)
- [Discharge Planning](#)
- [Quality Measures](#)
- [References](#)
- [Footnotes](#)
- [Codes](#)

Clinical Indications for Admission to Recovery Facility

[Return to top of Aortofemoral Bypass - RFC](#)

- Skilled nursing facility (SNF) or subacute facility care is/was needed for appropriate care of patient because of **ALL** of the following:
 - There are no acute hospital care needs.(1)
 - The patient has [Intense and complex care needs](#) that make recovery facility care safer and more practical than attempting care at a lower level and **ALL** of the following^{[A][B](4)}:
 - These care needs include the multiple components of care that are delivered by skilled professionals at a recovery facility.^{[C][D][E]}
 - There is a plan to provide **ALL** of the following⁽¹⁴⁾:
 - Care plan management and evaluation to meet patient needs, achieve treatment goals, and ensure medical safety⁽¹⁵⁾
 - Observation and assessment of patient's changing condition to evaluate need for treatment modification or for additional procedures until condition stabilized
 - Education services to teach patient self-maintenance or to teach caregiver patient care⁽¹⁶⁾⁽¹⁷⁾
 - [Skilled care](#) treatments daily (or more frequent), including **1 or more** of the following:
 - Nursing treatments needed for **1 or more** of the following⁽¹⁸⁾:
 - Intravenous (IV) infusion, IV injection, or intramuscular (IM) injection⁽¹⁾⁽¹⁶⁾
 - Tube feeding (ie, gastrostomy tubes, jejunostomy tubes, percutaneous endoscopic gastrostomy (PEG) tubes, nasogastric tubes) required because patient needs feeding to supply at least 26% of daily calories and at least 501 mL of daily fluids⁽¹⁶⁾⁽¹⁹⁾⁽²⁰⁾
 - Wound management that requires dressing changes with prescription medication or clean technique and treatment for **1 or more** of the following⁽¹⁾:
 - Surgical wound complications
 - Treatment with any stage III or IV pressure injury⁽¹⁶⁾
 - Treatment with 2 or more ulcers, including venous ulcers, arterial ulcers, or stage II pressure